# 

# Life Cycle Assessment: Results

The following supplementary LCA results are to be read alongside the complete ROCKWOOL<sup>®</sup> Environmental Product Declaration, attached.

## ROCKWOOL<sup>®</sup> stone wool product: ROCKLAP H&

ROCKLAP H&V Section

The results are for: 1 linear metre of product, Inner diameter of pipe section: with a thickness of 25 mm. 17 mm

#### Limitations

Conservative choices are made in the LCA as described in the ROCKWOOL<sup>®</sup> Group LCA rules. Therefore, the results can be considered to be conservative and worst case.

### Description of the system boundaries (x=included, MNA = Module not assessed)

	Prod	luct st	age	Constr instal sta	lation				Use stage	e			Er	nd-of-l	ife sta	ge	and loads beyond the system boundarie
	Kaw materials	Transport	Manufacturing	Transport	Assembly	Use	Maintenance	Repair	Replacement	Refurbishment	Operational energy use	Operational water use	De-construction demolition	Transport	Waste processing	Disposal	Reuse- Recovery- Recycling- potential
A	<b>\1</b>	A2	A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
	х	х	х	х	х	х	MNA	MNA	MNA	MNA	MNA	MNA	х	х	х	х	х

## Environmental impact

Parameter	Unit	A1-3	A4	A5	B1	C2	C4	D		
Global warming	kg $CO_2$ eqv	6.8E-01	7.5E-02	7.6E-02	0	1.7E-03	7.2E-03	-1.9E-02		
The global warming potential of a gas refers to the total contribution to global warming resulting from the emission of one unit of that gas relative to one unit of the reference gas, carbon dioxide, which is assigned a value of 1.										
Ozone depletion	kg CFC11 eqv	1.1E-09	1.2E-17	1.2E-10	0	5.6E-16	3.1E-14	-1.1E-15		
	fozone is caused	by the breakdo	h shields the earth wn of certain chlo hen they reach the molecules.	rine and/or bro	mine co	ntaining comp	ounds			
Acidification Acid depositions hav sources for emission		bstances are ag		il fuel combust				-6.3E-05		
Eutrophication	kg PO4 <sup>3-</sup> eqv	4.9E-04	1.2E-05	1.5E-05	0	7.7E-07	5.4E-06	-8.7E-06		
Excessive enrichme	ent of waters and	continental sur	faces with nutrien	its, and the ass	ociated a	adverse biolog	ical effects.			
Photochemical ozone creation	kg Ethene eqv	1.7E-04	-2.5E-07	4.0E-06	4.1E-11	-7.3E-07	3.6E-06	-7.2E-06		
Chemical reactions brought about by the light energy of the sun. The reaction of nitrogen oxides with hydrocarbons in the presence of sunlight to form ozone is an example of a photochemical reaction.										
Depletion abiotic resources -elements	kg Sb eqv	9.8E-06	6.3E-09	1.6E-09	0	1.3E-10	2.7E-09	-4.2E-09		
Depletion abiotic resources fuels	MJ	8.5E+00	1.0E+00	1.3E-01	0	2.3E-02	1.0E-01	-4.7E-01		
Consumpt	tion of non-renew	able resources,	thereby lowering	their availabili	ty for fut	ure generation	ns.			

# 

### Resource use

Parameter	Unit	A1-3	A4	A5	B1	C2	C4	D
Use of renewable primary energy excluding renewable primary energy resources used as raw materials	MJ	2.2E+00	5.7E-01	7.1E-01	0	1.4E-03	1.3E-02	-2.6E-01
Use of renewable primary energy resources used as raw materials	MJ	8.6E-01	0.0E+00	-6.5E-01	0	0.0E+00	0.0E+00	0.0E+00
Total use of renewable primary energy resources	MJ	3.1E+00	5.8E-02	5.1E-02	0	1.4E-03	1.3E-02	-2.6E-01
Use of non-renewable primary energy excluding non-renewable primary energy resources used as raw materials	MJ	9.1E+00	1.0E+00	1.5E-01	0	2.3E-02	1.0E-01	-4.9E-01
Use of non-renewable primary energy resources used as raw materials	MJ	1.1E+00	0.0E+00	-2.6E-03	0	0.0E+00	0.0E+00	0.0E+00
Total use of non-renewable primary energy resources	MJ	1.0E+01	1.0E+00	1.4E-01	0	2.3E-02	1.0E-01	-4.9E-01
Use of secondary materials	kg	0.0E+00	n/a	0.0E+00	n/a	n/a	n/a	n/a
Use of renewable secondary fuels	MJ	*	*	*	*	*	*	*
Use of non-renewable secondary fuels	MJ	*	*	*	*	*	*	*
Net use of fresh water	m <sup>3</sup>	3.9E-03	6.7E-05	1.9E-04	0	1.1E-06	2.4E-05	-1.7E-04

\* There are no renewable and no non-renewable secondary fuels used in A3. The minor use of secondary fuels as part of the background datasets is not accounted for.

### Waste categories

Parameter	Unit	A1-3	A4	A5	B1	C2	C4	D
Hazardous waste disposed	kg	2.2E-06	4.8E-08	3.2E-09	0	5.7E-09	9.3E-09	-8.6E-10
Non-hazardous waste disposed	kg	6.2E-02	1.6E-04	1.1E-02	0	4.2E-06	5.0E-01	-1.3E-03
Radioactive waste disposed*	kg	3.2E-04	1.3E-06	3.2E-06	0	2.9E-08	1.3E-06	-6.9E-07

\* There is never radioactive waste from a ROCKWOOL plant (A3), but there might be small amounts associated with the secondary LCI datasets used for the upstream chain (A1 & A2), which are taken into account here.

### Output flows

Parameter	Unit	A1-3	A4	A5	B1	C2	C4	D
Component for re-use	kg	1.06E-07	n/a	3.15E-09	n/a	n/a	n/a	n/a
Materials for recycling	kg	1.86E-02	n/a	n/a	n/a	n/a	n/a	n/a
Materials for energy recovery	kg	2.14E-05	n/a	n/a	n/a	n/a	n/a	n/a

